



ELSEVIER

Mechanisms of Development 58 (1996) 237



Volume contents

Volume 58 (1996)

Issue contents

| | | | |
|---|-----|---|-----|
| M. Hoch, M.J. Pankratz Control of gut development by <i>fork head</i> and cell signaling molecules in <i>Drosophila</i> | 1 | K.R. Kao, A. Lockwood Negative regulation of dorsal patterning in early embryos by overexpression of <i>XrelA</i> , a <i>Xenopus</i> homologue of NF- κ B | 129 |
| J. Klingensmith, Y. Yang, J.D. Axelrod, D.R. Beier, N. Perrimon, D.J. Sussman Conservation of <i>dishevelled</i> structure and function between flies and mice: isolation and characterization of <i>Dvl2</i> | 3 | P. Bouillet, M. Oulad-Abdelghani, S.J. Ward, S. Bronner, P. Chambon, P. Dollé A new mouse member of the <i>Wnt</i> gene family, <i>mWnt-8</i> , is expressed during early embryogenesis and is ectopically induced by retinoic acid | 141 |
| A.-L. Gustafson, M. Donovan, E. Annerwall, L. Dencker, U. Eriksson Nuclear import of cellular retinoic acid-binding protein type I in mouse embryonic cells | 15 | N.M.T. van der Lugt, M. Alkema, A. Berns, J. Deschamps The <i>Polycomb</i> -group homolog <i>Bmi-1</i> is a regulator of murine <i>Hox</i> gene expression | 153 |
| D.I. Strutt, M. Mlodzik The regulation of <i>hedgehog</i> and <i>decapentaplegic</i> during <i>Drosophila</i> eye imaginal disc development | 27 | A. Mallamaci, E. Di Blas, P. Briata, E. Boncinelli, G. Corte OTX2 homeoprotein in the developing central nervous system and migratory cells of the olfactory area | 165 |
| E. Bloch-Gallego, S. Millet, R.-M. Alvarado-Mallart Further observations on the susceptibility of diencephalic prosomeres to <i>En-2</i> induction and on the resulting histogenetic capabilities | 39 | C. Borchiellini, J. Coulon, Y. Le Parco The function of type IV collagen during <i>Drosophila</i> muscle development | 179 |
| H. Xiao, L.A. Hrdlicka, J.R. Nambu Alternate functions of the <i>single-minded</i> and <i>rhomboid</i> genes in development of the <i>Drosophila</i> ventral neuroectoderm | 51 | S. Lu, C.S. Shashikant, F.H. Ruddle Separate <i>cis</i> -acting elements determine the expression of mouse <i>Dbx</i> gene in multiple spatial domains of the central nervous system | 193 |
| M. Nemer, E.W. Stuebing <i>WEE1</i> -like CDK tyrosine kinase mRNA level is regulated temporally and spatially in sea urchin embryos | 65 | W. Wang, X. Chen, H. Xu, T. Lufkin <i>Msx3</i> : a novel murine homologue of the <i>Drosophila</i> <i>msh</i> homeobox gene restricted to the dorsal embryonic central nervous system | 203 |
| J. den Hertog, J. Overvoorde, S.W. de Laat Expression of receptor protein-tyrosine phosphatase α mRNA and protein during mouse embryogenesis | 75 | M. D'Alessio, M. Frasch <i>msh</i> may play a conserved role in dorsoventral patterning of the neuroectoderm and mesoderm | 217 |
| M. Baltzinger, F. Relaix, P. Remy Transcription of <i>XLPOU3</i> , a brain-specific gene, during <i>Xenopus laevis</i> early embryogenesis | 89 | Author index | 233 |
| B. Lefevre, F. Crossin, J. Fontaine-Pérus, E. Bandman, M.-F. Gardahaut Innervation regulates myosin heavy chain isoform expression in developing skeletal muscle fibers | 103 | Subject index | 237 |
| | 115 | Volume contents | 239 |